

EMG evaluation of hip adduction exercises for soccer players: implications for exercise selection in prevention and treatment of groin injuries

Andreas Serner,^{1,2} Markus Due Jakobsen,³ Lars Louis Andersen,³ Per Hölmich,^{1,2} Emil Sundstrup,³ Kristian Thorborg¹

¹Arthroscopic Centre Amager, Copenhagen University Hospital, Copenhagen, Denmark

²Aspetar Sports Groin Pain Centre, Aspetar, Qatar Orthopaedic and Sports Medicine Hospital, Doha, Qatar

³National Research Centre for the Working Environment, Copenhagen, Denmark

Correspondence to

Andreas Serner, Arthroscopic Centre Amager, Copenhagen University Hospital, Italiensvej 1, Copenhagen 2300, Denmark; andreasserner@hotmail.com

Accepted 19 February 2013

ABSTRACT

Introduction Exercise programmes are used in the prevention and treatment of adductor-related groin injuries in soccer; however, there is a lack of knowledge concerning the intensity of frequently used exercises.

Objective Primarily to investigate muscle activity of adductor longus during six traditional and two new hip adduction exercises. Additionally, to analyse muscle activation of gluteals and abdominals.

Materials and methods 40 healthy male elite soccer players, training >5 h a week, participated in the study. Muscle activity using surface electromyography (sEMG) was measured bilaterally for the adductor longus during eight hip adduction strengthening exercises and peak EMG was normalised (nEMG) using an isometric maximal voluntary contraction (MVC) as reference. Furthermore, muscle activation of the gluteus medius, rectus abdominis and the external abdominal obliques was analysed during the exercises.

Results There were large differences in peak nEMG of the adductor longus between the exercises, with values ranging from 14% to 108% nEMG ($p < 0.0001$). There was a significant difference between legs in three of the eight exercises (35–48%, $p < 0.0001$). The peak nEMG results for the gluteals and the abdominals showed relatively low values (5–48% nEMG, $p < 0.001$).

Conclusions Specific hip adduction exercises can be graded by exercise intensity providing athletes and therapists with the knowledge to select appropriate exercises during different phases of prevention and treatment of groin injuries. The Copenhagen Adduction and the hip adduction with an elastic band are dynamic high-intensity exercises, which can easily be performed at any training facility and could therefore be relevant to include in future prevention and treatment programmes.

INTRODUCTION

Groin injuries are one of the most common types of injuries in soccer accounting for 8–18% of all injuries.^{1–7} It is described that the typical groin injury in soccer is primarily related to the hip adductors, which account for 69% of all groin injuries.⁸

There is consistent evidence that previous groin injury increases the risk for subsequent groin injury.^{1 5 9 10} A common assumption is that strength deficits due to inadequate rehabilitation is a possible reason for previous injuries being a risk factor.^{1 5 9} Additionally, reduced hip adductor strength has been found to be a risk factor for subsequent groin injury.^{9 11} Therefore, hip adduction strengthening exercises seem to have a large

potential in both the prevention and treatment of groin injuries.

Currently, no studies have been able to demonstrate a reduction in the number of groin injuries in soccer.^{12–14} Various explanations for this have been proposed, such as insufficient compliance,^{12–14} over-optimistic effect sizes¹² and inadequate exercise intensity.¹² Physical training has proven effective in the treatment of long-standing adductor-related groin pain,¹⁵ which is supported by other studies showing favourable outcomes of exercise interventions.¹⁶ None of the aforementioned studies,^{12–15 17–20} however, account for the intensity of the included exercises, that is, relative load, perceived exertion or muscle activity. Consequently, no information exists on the specific intensity of exercises in the treatment of groin pain in athletes. In this regard, electromyography is a pertinent method of evaluating exercise intensity.^{21–24} The results of the present study could therefore be used to optimise future training programmes by a greater understanding of exercise intensity when selecting exercises to match different contextual situations and physiological demands in prevention and treatment of adductor-related groin injuries.

The primary purpose of this study was therefore to examine the muscle activity, defined as the peak normalised EMG signal (nEMG) of the adductor longus during six traditional and two new hip adduction strengthening exercises. Additionally, muscle activity of the gluteus medius, rectus abdominis and external abdominal oblique was evaluated.

MATERIALS AND METHODS

Subjects

Forty healthy male elite soccer players older than 18 years participated in the study (21.4 ± 3.3 years, 182.4 ± 7.9 cm, 77.8 ± 7.9 kg). The participants were recruited on a voluntary basis through coaches and physiotherapists from teams from the Danish Senior 1st Division, 2nd Division and the Under-19 League. All participants were training >5 h per week with 1–2 weekly matches. Players were eligible for inclusion if they did not have current hip or groin symptoms and/or injury or other injuries which could be presumed to influence the execution of the tests. As a large proportion of soccer players experience mild hip and groin symptoms and disability, that may not be perceived as problematic by the player, a score higher than 80 out of 100 in the subscale ‘function, sports and recreational activities’ from the Copenhagen Hip and Groin Outcome Score,²⁵ HAGOS (Sport), was required. Furthermore, we

To cite: Serner A, Jakobsen MD, Andersen LL, et al. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2012-091746

decided to exclude players if they reported pain >2 on a numeric pain rating scale in the Adductor Functional Pain Test²⁶ during maximal voluntary contractions (MVCs) and during the performance of the eight exercises. Inclusion of participants ended when the desired sample size was reached. No exercise or training was allowed on the same day before testing and all participants gave written informed consent according to the Helsinki Declaration before testing was initiated. The study was approved by the Danish National Committee on Health Research Ethics (H-3-2011-145).

Testing procedure

The participants attended two separate test sessions. On the first session, they performed a 10 min standardised warm-up and were subsequently familiarised with all the exercises. The second session was conducted with a minimum of 3 days' interval to avoid delayed onset muscle soreness (DOMS). After the warm-up, the players performed the MVC tests followed by the eight exercises.

Hip adduction exercises

The present study included six traditional hip adduction exercises^{12 15 27 28} and two new exercises which were considered to activate the hip adductor muscle group with a different intensity (figure 1). We wanted to introduce exercises that we thought would represent both high and low intensities, as this can be useful in clinical situations, such as acute versus chronic conditions, where muscle-tendinous structures need to be targeted with different intensities. Only exercises where the hip adductors were used as a prime mover in hip adduction were included to ensure similar muscle actions in our comparisons. The preferred kicking leg was defined as the dominant leg and was used as the training leg when performing the exercises.

The exercises were performed in a randomised order and a minimum of 30 s break was held between the exercises.²⁹ EMG increases due to fatigue with consecutive repetitions within a set performed to failure,³⁰ but not between consecutive sets with rest periods in between.³¹ To avoid fatigue, the subjects therefore performed two separate sets and the dynamic exercises were performed with three repetitions and one repetition for the isometric exercises in each set. All exercises were performed for 6 s per repetition: 3 s concentric and a 3 s eccentric contraction for the dynamic exercises, with a gradual increase and decrease of tension for the six second isometric exercises.

For the two exercises that can be adjusted using an external load, the hip adductor machine and the hip adduction with an elastic band, a relative load of 10 repetition maximum (RM), was chosen. The load on the machine and the resistance of the elastic bands (number and strength of the elastic bands and distance from the fixation point) were individually determined on the first session. The participants subjectively evaluated the exercises after 5–6 repetitions and adjusted the load accordingly. Ten repetitions were performed and the relevant load was noted if participants considered the load equivalent to 10 RM. A load of 10 RM is described as heavy resistance training³² and has been used previously in hip adduction strengthening,²⁷ and therefore it is considered as an appropriate reference for exercise comparisons.

The eight exercises

(a) *Isometric adduction—with a ball between the ankles* (IBA): Isometric hip adduction against a football placed between the feet inferior to the medial malleoli when lying supine. The knees are kept straight and feet vertical.

The feet are pressed against the ball as hard as possible. Exercise used in refs. 12 15 28.

(b) *Isometric adduction—with a ball between the knees* (IBK): Isometric hip adduction against a football placed between the knees when lying supine with hips and knees flexed and feet flat on the surface. The first toe is pointed straight forward and the knees are pressed against the ball as hard as possible. Exercise used in refs. 12 15 28.

(c) *Side-lying hip adduction* (SLA): The player is lying on the side of the dominant leg with the dominant leg straight and the non-dominant leg flexed 90° in the hip and knee. Maximal hip adduction of the dominant leg is performed




Figure 1 The eight included hip adduction exercises: (a) isometric adduction with a ball between the ankles; (b) isometric adduction with a ball between the knees; (c) side-lying adduction; (d) sliding hip adduction/adduction exercise; (e) hip adduction with an elastic band; (f) hip adductor machine; (g) supine bilateral hip adduction; (h) Copenhagen adduction.

keeping the knee straight and foot horizontal. Exercise used in ref. 15.

- (d) *Sliding hip abduction/adduction* (SLIDE): Standing with the dominant leg on a wash cloth with hips and knees straight and hands placed laterally on the trunk. The dominant leg is abducted, so the centre of gravity moves laterally as much as controllably possible. The dominant leg is then adducted, sliding back to the starting position. Exercise modified from refs. 15 28.
- (e) *Hip adduction with an elastic band* (EB): Standing with the hip of the dominant leg positioned in maximal abduction. An elastic band (Thera-Band) is attached to a fixation point and around the ankle of the dominant leg keeping a slight tension in the elastic band from the beginning. The upper body is kept straight and the player holds on to a stable support. The dominant leg is adducted until there is a foot width between the feet and the dominant leg is approximately half a foot length behind the foot of the non-dominant leg. Exercise modified from refs. 15 27 28.
- (f) *Hip adductor machine* (AM): Sitting in a hip adduction machine, 'Adductor' (TechnoGym, Bologna, Italy), with a slight knee flexion holding on to the handles. The legs are then moved into maximal hip abduction and back. Exercise used in ref. 28.
- (g) *Supine bilateral hip adduction* (SBA): Supine with the hips in 90° flexion and the knees in 45° flexion. The legs are moved into maximal hip abduction and back keeping the flexion in the hips and knees constant. New suggested low-intensity exercise.
- (h) *Copenhagen Adduction* (CA): A partner exercise where the player is lying on the side of the non-dominant leg with one forearm as support on the floor and the other arm placed along the body. The dominant leg is held in approximately the height of the hip of the partner, who is holding the leg with one hand supporting the ankle and the other supporting the knee. The player then raises the body from the floor and the non-dominant leg is adducted so that the feet touch each other and the body is in a straight line. The body is then lowered halfway to the ground while the foot of the non-dominant leg is lowered so that it just touches the floor without using it for support. New suggested high-intensity exercise.

Electromyography

Bipolar disposable silver/silver chloride (Ag/AgCl) surface electrodes (White Sensor, Ambu A/S, Ballerup, Denmark) were used bilaterally to collect data from the adductor longus, gluteus medius, rectus abdominis and the external abdominal oblique. The electromyography (EMG) activity was sampled at 1024 Hz using two sets (4 channels each) of the Nexus 10 data loggers (Mind Media, The Netherlands) and digitally converted at a 24 bit resolution using integrated A/D-converters. To reduce impedance, the area where the electrodes were placed was shaved and cleansed with an abrasive gel (Acqua gel, Meditec, Parma, Italy).

Electrode placement

The electrodes were placed parallel to the muscle fibres with an inter-electrode distance of 2 cm to avoid unstable recordings due to tendon and motor endplate effects, as recommended by the SENIAM project.³³ To confirm the correct placement of electrodes, a manual muscle test was used and the EMG signal was checked on a laptop via Bluetooth. The same tester

conducted preparation and control for all participants, and electrodes were placed in the following way:

Adductor longus: medially on the thigh equivalent to the proximal third of the distance from the pubic tubercle to the insertion on the femur.^{34 35} *Gluteus medius*: half the distance between the iliac crest and the major trochanter anterosuperiorly from the Gluteus Maximus.^{33–36} *Rectus abdominis*: half the distance between the umbilicus and the pubic symphysis³⁷ and centred on the muscle belly, that is, 2–4 cm laterally from the midline depending on the size of the participant. *External abdominal obliques*: directly below the most inferior point of the costae in direction towards the opposite pubic tubercle.^{38 39} *Reference electrodes*: on the medial surface of the tibia. All electrodes were fixated with tape (Fixomull Stretch, BSNmedical, Germany).

Reference contractions

The participants performed two MVCs in six different test positions and in a randomised order to determine the maximal muscle activity. The trial with the highest EMG value was used as the reference contraction, that is, 100%, to normalise the peak EMG recordings from the exercises. MVCs lasted 5 s each and participants were given at least 30 s rest between each test.²⁹ A standardised command was used during tests; 'go ahead, push, push, push, push and relax', which has been used previously in the evaluation of maximal strength recordings.⁴⁰ MVCs were conducted in the following way:

Adductor longus MVC: a bilateral hip adduction was performed supine with the hips and knees straight and a ball placed between the knees. *Gluteus medius MVC*: a maximal hip abduction was performed side-lying with knees straight and the tested leg in 25° of hip abduction.^{36 41} A strap belt around the knees just proximal of the lateral femur epicondyle was used as it has been shown to increase the reliability of maximal hip abduction measurements.⁴² *Rectus abdominis and external abdominal obliques*: Two test positions were performed to ensure the highest magnitude of muscle activity.⁴³ A straight abdominal flexion in an 'Abdominal Crunch R.O.M' machine (TechnoGym, Bologna, Italy) and a trunk rotation in a 'Rotary Torso' machine (TechnoGym) to each side. Both MVCs were performed isometrically with the resistance and movement of the machines locked.

Data extraction

All raw EMG signals were digitally filtered by a Butterworth fourth-order high-pass filter (10 Hz cut-off frequency). Raw EMG was checked visually on a computer and processed to ensure correct signal registration. For each individual muscle, a moving root mean square (RMS) EMG (500 ms time constant) was used to identify peak EMG within each contraction (repetition) and normalised to the maximal moving RMS EMG (500 ms time constant) obtained during MVC.^{22 30 31} Reported peak nEMG for each exercise was calculated using the average of the peak nEMG from all repetitions, that is, six repetitions for the dynamic exercises and two repetitions for the isometric exercises.

Statistical methods

Intraclass correlation coefficient (ICC) model 3 was used to determine the intrasession reliability of the repetitions (six dynamic and two isometric) using the peak nEMG from all subjects in each exercise. As a reference, ICC values >0.75 were considered excellent and 0.40–0.75 fair to good.⁴⁴ A repeated-measure one-way analysis of variance (ANOVA) was used for the evaluation of peak nEMG for each muscle with nEMG as

the dependent variable and *exercise* as the independent variable. A post hoc analysis was made to compare the exercises and a Bonferroni correction was made to account for the risk of mass significance, that is, all *p* values presented have been multiplied by the relevant number of pairwise comparisons. When comparing differences between the adductor muscle activity of the two legs, a two-way ANOVA was used to test for muscle by exercise interaction, with *nEMG* as the dependent variable and *muscle* (dominant and non-dominant adductor longus) and *exercise* as independent variables. All *nEMG* values are reported as least square mean \pm SE and the level of significance was set at $p < 0.05$.

Sample size calculation

The SD in *nEMG* for leg exercises has been shown to be approximately 15%.²⁴ The minimal clinically relevant difference between exercises has been suggested to be 10% *nEMG*.^{22–45} With a power of 80% (β 0.80) and a significance level of 5% ($\alpha=0.05$), the minimum number of participants was calculated to be 20. To account for multiple comparisons and missing data (eg, due to electrode errors etc.), we decided to include 40 participants.

RESULTS

Subjects

Data from two participants were missing due to broken reference electrodes. Therefore, the analysis included 38 participants (table 1). Full datasets were registered for 35 participants and data from three participants had a total of 18 single contraction measurements showing uncorrectable registration errors, which led to their exclusion from the respective analyses. Analysis of peak EMG values was therefore conducted to account for the missing values.

Reliability of repetitions

For the majority of the exercises, excellent ICC values were found between the repetitions for adductor longus (ICC 3=0.81–0.97). The ICC values for the dominant adductor longus in the isometric adduction with the ball between the knees and between the ankles were fair to good with an ICC of 0.59 and 0.74, respectively.

Peak muscle activity

We found a significant exercise by muscle interaction (ANOVA: $p < 0.0001$) for the adductor longus, as well as a significant difference between the exercises for the gluteus medius, rectus abdominis and the external abdominal oblique muscles (ANOVA: $p < 0.0001$). See table 2 for specifications.

Post hoc analysis

The post hoc analysis of adductor longus activation (table 3) showed that the supine bilateral hip adduction had significantly lower values than all other exercises ($p < 0.01$). The side-lying hip adduction was significantly higher than the supine bilateral hip adduction and significantly lower than all the other exercises

($p < 0.01$). Furthermore, there was a significant difference from the three exercises with the highest *nEMG* activity and the three lowest ranked exercises ($p < 0.05$) (table 3).

Between-leg difference

Three of the exercises, the Copenhagen Adduction, the hip adduction with an elastic band and the side-lying hip adduction, had significantly higher adductor longus activation in the dominant leg (training leg) compared to the non-dominant leg (non-training leg), 35–48%, $p < 0.0001$ (figure 2). Consequently, these exercises can be considered asymmetric regarding intensity.

DISCUSSION

The main purpose of this study was to evaluate the exercise intensity using sEMG to measure muscle activity during the hip adduction exercises frequently used in the prevention and treatment of adductor-related groin injuries. The findings of this study show that there is a significant difference in muscle activation between the included exercises.

This evaluation of exercise intensity can help guide clinical decision making by an increased understanding of ways to progress strength training exercises in the prevention and treatment of adductor-related injuries. The results show that seven exercises can be considered strength training, as they elicit an intensity higher than 60% of maximum, which is documented to be conducive of muscle strength gains.⁴⁶ Furthermore, it could be assumed that it is possible to regulate the intensity of some of the exercises by changing the RM, both to a higher or lower relative load or by pressing submaximally on the ball in the isometric exercises.

There was no significant difference between the five exercises with the highest *nEMG* values. These exercises are all performed in different test positions and with different kinds of equipment (elastic band, machine, wash cloth, ball or without anything), allowing the clinician to choose the exercises which fit the players' situation and preferences best. This also shows promising possibilities for implementing strength training on the pitch for football players, as the Copenhagen Adduction exercise, the hip adduction with an elastic band and the isometric hip adduction with a ball between the knees all can be performed at any training facility with a minimum of equipment needed. The optimal number of sets, repetitions and frequency of the exercises is, however, beyond the scope of this study.

There is limited research investigating adductor longus muscle activity during exercise. A review investigating the free barbell squat finds that a greater hip external rotation position, a wide stance of the feet, as well as an increased load will increase hip adduction activation during the exercise.⁴⁷ The highest *nEMG* value found for the hip adductors was, however, only 23.1%.⁴⁸ A different study, which investigated adductor longus activation during a single-leg squat, a lunge and a step up-and-over exercise, showed peak *nEMG* values at 15–19%, 8–21% and 15–22% *nEMG*, respectively.⁴⁹ The relatively low *nEMG* results found in these exercises indicate that they probably do not sufficiently activate the adductors to achieve strength increases and hypertrophy. Consequently, seven exercises from the present study appear to be superior in that respect, as they all focus primarily on the hip adduction movement, thereby reaching greater levels of muscle activity.

The adductor muscle group's main action is described to be adduction of the thigh in the open kinetic chain and stabilisation of the lower extremity in the closed kinetic chain.⁵⁰ The variability of soccer play stresses the adductor muscle group in different situations with both the high running loads with

Table 1 Baseline data for 38 participants

Age	21.4 \pm 3.3 years
Height	182.4 \pm 7.9 cm
Weight	77.8 \pm 7.9 kg
Adductor functional pain test	0 \pm 0
HAGOS (Sport)	95.8 \pm 5.7 points

Table 2 Peak normalised EMG measurements for all exercises and muscles

Exercise	Adductor longus		Gluteus medius		External abdominal oblique		Rectus abdominis	
	Dominant side	Non-dominant side	Dominant side	Non-dominant side	Dominant side	Non-dominant side	Dominant side	Non-dominant side
1. Isometric adduction with a ball between the knees	108±6	102±6	10±3	25±3	13±3	11±3	8±3	8±3
2. Copenhagen adduction	108±5	69±6	20±3	48±3	29±3	36±3	19±3	40±3
3. Hip adduction with an Elastic band	103±6	70±6	18±3	18±3	18±3	10±3	9±3	6±3
4. Hip adductor machine	99±6	95±6	12±3	15±3	15±3	10±3	11±3	8±3
5. Sliding hip abduction/adduction exercise	98±6	89±6	36±3	45±3	14±3	11±3	7±3	5±3
6. Isometric adduction with a ball between the ankles	86±6	80±6	13±3	18±3	18±3	18±3	21±3	17±3
7. Side-lying hip adduction	64±6	17±6	22±3	14±3	12±3	21±3	5±3	13±3
8. Supine hip adduction	14±6	15±6	16±3	21±3	35±3	31±3	36±3	30±3

Values reported as mean peak normalised EMG± standard error in % of the MCV and listed in the order of adductor longus activation. Dominant side=side of the preferred kicking leg.

high intensity turns and side-cutting manoeuvres, as well as the repetitive action of the soccer kick being stressful components.^{51 52}

These movements can result in large amounts of stretching of highly activated muscles (lengthening contractions), which are associated with muscle injuries.^{53–56} Additionally, these movements create large stabilisation requirements around the hip, which if inadequate might also be a risk factor for injury. This was demonstrated in a study of ice hockey players, where a muscular imbalance with a preseason hip adduction/abduction strength ratio less than 80% increased the risk of sustaining an adductor muscle strain in the following season by 17 times,⁵⁷ emphasising the importance of ensuring adequate strength of the adductors.

The included dynamic exercises appear to be superior to the isometric exercises as they include the element of lengthening contractions, thereby preparing the muscle-tendinous structures for the requirements of soccer. Exercises where the adductor longus muscle is working eccentrically seem important to consider, especially in prevention, as this resembles the situation where the muscle-tendinous structures are at the highest risk of

injury. Three of the exercises, the Copenhagen Adduction, the hip adduction with an elastic band and the hip adductor machine, all include eccentric contractions and a high nEMG activity of the adductor longus. They might therefore be recommended as the potentially best exercises in the prevention of adductor-related groin injuries in soccer players, although future studies need to confirm a possible preventive effect. These findings are supported by a study showing that an 8-week progressive strengthening programme using a dynamic hip adduction exercise with an elastic band increased eccentric hip adduction strength by 13%.²⁷

A substantiating argument for the potential of dynamic high-intensity exercises for soccer players is that an exercise, matching the proposed injury mechanisms and with high muscle activity, has been able to significantly prevent another muscle-tendinous injury.⁵⁸ This study includes the Nordic Hamstring exercise, which has shown a biceps femoris peak nEMG of 91%,⁴⁵ and is performed eccentrically to match the deceleration phase of the leg during running/sprinting, where hamstring injuries are described to occur.^{59 60}

Table 3 Pairwise comparisons adductor longus (dominant leg)

Exercise	2. Copenhagen adduction (108%)	3. Hip adduction with an elastic band (103%)	4. Hip adductor machine (99%)	5. Sliding hip abduction/adduction (98%)	6. Isometric adduction with a ball between the ankles (86%)	7. Side-lying hip adduction (64%)	8. Supine bilateral hip adduction (14%)
1. Isometric adduction with a ball between the knees (108%)	0±5	5±5	9±5	10±5	22±5**	44±5**	94±5**
2. Copenhagen adduction (108%)		5±5	9±5	10±5	22±5**	44±5**	94±5**
3. Hip adduction with an elastic band (103%)			4±5	5±5	17±5*	39±5**	89±5**
4. Hip adductor machine (99%)				1±5	13±5	35±5**	85±5**
5. Sliding hip abduction/adduction (98%)					11±5	34±5**	83±5**
6. Isometric adduction with a ball between the ankles (86%)						23±5**	72±5**
7. Side-lying hip adduction (64%)							49±5**

*p<0.05 and **p<0.01 (Bonferroni corrected).

Values are reported as the difference of least square means in % nEMG± SE. Exercises are listed according to the level of peak nEMG of the dominant adductor longus with peak nEMG stated in brackets. nEMG=normalised electromyography.




Figure 2 Mean peak nEMG for the dominant and the non-dominant adductor longus. nEMG=Normalised electromyography; Dominant leg=The preferred kicking leg. * Denotes a significant difference between the dominant and the non-dominant leg. IBK, isometric adduction with a ball between the knees; CA, Copenhagen adduction; EB, hip adduction with an elastic band; AM, hip adductor machine, SLIDE, sliding hip abduction/adduction exercise; IBA, isometric adduction with a ball between the ankles; SLA, side-lying adduction; SBA, supine bilateral hip adduction.

Our findings show a significant asymmetry in muscle activation between legs in three exercises. This information can assist in the selection of exercises, for example, if a differentiated training load between legs is desired. We found no significant differences in muscle activation between legs in the symmetrically performed exercises. Previous research has documented strength asymmetries between legs in elite soccer players^{61 62}; however, there is no indication that this has any influence on our EMG findings in the symmetrical exercises.

The muscle activity of the secondary muscles (abdominals and gluteals) all show values of less than 49% nEMG. Consequently, the included exercises are not ideal for strength training of these specific muscle groups in healthy athletes. In comparison, other studies have found values at 16–32% nEMG and 39–64% for the external abdominal oblique and rectus abdominis, respectively, during abdominal crunch exercises.^{63–66} Both considerable movement and stabilisation of the trunk are required in soccer, especially during kicking,⁶⁷ and recurrent stress on the abdominal muscles during soccer has been suggested to be involved in the aetiology of groin pain.⁶⁸ Therefore, choosing exercises with a high abdominal activation also seems relevant in relation to groin injuries. To target the abdominal muscle-tendinous structures to some extent, through the exercises included in the present study, the Copenhagen Adduction exercise seems preferred, as it shows the highest values for the abdominal muscles reaching 36% and 40% peak nEMG for the external abdominal oblique and rectus abdominis on the non-dominant side, respectively.

Limitations

The present study was conducted on male soccer players only and therefore limits the generalisation to this group. Consequently, we cannot conclude that the results are applicable to females, other types of athletes, players with pathologies or the general untrained population.

Surface EMG is subject to various limitations, as the EMG signal is influenced by factors such as thickness of tissue layers, possible

cross talk, electrode shifting, changes in muscle tissue conductivity, etc.^{69 70} To reduce variability as much as possible, the same tester conducted the preparation of all participants using a precisely described placement of electrodes. Additionally, this study utilised standardised reference contractions. Furthermore, a 500 ms RMS window was used in the analysis to minimise the influence of rapid changes in muscle activity during the repetitions.

The literature lacks consistency in the description of MVCs for the adductor longus.^{49 51 71 72} Consequently, the MVC test position was chosen because the majority of the exercises in the present study are performed with the hips and knees straight. We found that the MVC had a lower peak EMG than some of the exercises. Therefore, the results of the present study might not be reproducible when using a different MVC test for normalisation. Based on the results of this study, an isometric hip adduction with hip and knee flexion and a ball between the knees should be considered as an MVC in future studies concerning adductor longus EMG measurements.

Two ICC measurements for the dominant adductor longus showed less than excellent values. These were from the isometric exercises. We did not find any significant differences in peak nEMG between legs in the isometric exercises, and as the ICC values for the EMG activity in the non-dominant leg in the isometric tests were excellent (0.89), we do not consider the lower ICC values for the dominant leg to have a significant influence on our findings. This could be due to the large sample size reducing this possible measurement variation.

The use of a Bonferroni correction in the post-hoc analysis is a very conservative approach and increases the risk of type 2 errors, that is, false negatives. The post-hoc analyses show significant differences for the three exercises with the lowest peak nEMG, but not between the five exercises with the highest peak nEMG. These five exercises were, however, all within 10% nEMG, which was predetermined as clinically irrelevant. Thus, a possible type 2 error is not considered to be of clinical relevance.

CONCLUSION

Specific hip adduction exercises can be graded by exercise intensity, providing athletes and therapists with the knowledge to select appropriate exercises during different phases of prevention and treatment of groin injuries. The Copenhagen Adduction and the hip adduction with an elastic band are dynamic high intensity exercises, which can easily be performed at any training facility and therefore seem relevant to include in future prevention and treatment programmes.

What this study adds

- ▶ Specific hip adduction exercises display different levels of adductor longus muscle activity, indicating differences in exercise intensity between specific exercises.
- ▶ This enables a deliberate choice of exercises based on intensity, which is relevant in different phases of prevention and treatment of groin injuries.
- ▶ The Copenhagen Adduction exercise and the hip adduction exercise with an elastic band are dynamic high-intensity exercises with limited or no equipment. Therefore, they have great potential for on-field prevention and treatment programmes.

Acknowledgements The authors would like to thank all the players who took the time to participate in the study, as well as David Visvanathan and the team of physiotherapy students for their help with the practical work during the tests. Anders Vinther, PhD, is also acknowledged for his supporting input.

Contributors AS and KT contributed to the conception and design of the study. MDJ, ES and LLA assisted with the technical aspects of the exercise protocol. All authors contributed to the analysis and interpretation of the data. AS recruited all participants and supervised the exercise protocol. AS drafted the article and all authors revised it critically and gave final approval of the version submitted.

Competing interests None.

Patient consent Obtained.

Ethics approval The study was approved by the Danish National Committee on Health Research Ethics (H-3-2011-145).

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Arnason A, Sigurdsson SB, Gudmundsson A, *et al.* Risk factors for injuries in football. *Am J Sports Med* 2004;32:55–165.
- 2 Ekstrand J, Hilding J. The incidence and differential diagnosis of acute groin injuries in male soccer players. *Scand J Med Sci Sports* 1999;9:98–103.
- 3 Häggglund M, Waldén M, Ekstrand J. Injuries among male and female elite football players. *Scand J Med Sci Sports* 2009;19:819–27.
- 4 Ekstrand J, Häggglund M, Waldén M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med* 2011;45:553–8.
- 5 Häggglund M, Waldén M, Ekstrand J. Previous injury as a risk factor for injury in elite football: a prospective study over two consecutive seasons. *Br J Sports Med* 2006;40:767–72.
- 6 Waldén M, Häggglund M, Ekstrand J. Injuries in Swedish elite football—a prospective study on injury definitions, risk for injury and injury pattern during 2001. *Scand J Med Sci Sports* 2005;15:118–25.
- 7 Werner J, Häggglund M, Waldén M, *et al.* UEFA injury study: a prospective study of hip and groin injuries in professional football over seven consecutive seasons. *Br J Sports Med* 2009;43:1036–40.
- 8 Hölmich P. Long-standing groin pain in sportspeople falls into three primary patterns, a 'clinical entity' approach: a prospective study of 207 patients. *Br J Sports Med* 2007;41:247–52.
- 9 Engebretsen AH, Myklebust G, Holme I, *et al.* Intrinsic risk factors for groin injuries among male soccer players: a prospective cohort study. *Am J Sports Med* 2010;38:2051–7.
- 10 Steffen K, Myklebust G, Andersen TE, *et al.* Self-reported injury history and lower limb function as risk factors for injuries in female youth soccer. *Am J Sports Med* 2008;36:700–8.
- 11 Crow JF, Pearce AJ, Veale JP, *et al.* Hip adductor muscle strength is reduced preceding and during the onset of groin pain in elite junior Australian football players. *J Sci Med Sport* 2010;13:202–4.
- 12 Hölmich P, Larsen K, Krosgaard K, *et al.* Exercise program for prevention of groin pain in football players: a cluster-randomized trial. *Scand J Med Sci Sports* 2010;20:814–21.
- 13 Engebretsen AH, Myklebust G, Holme I, *et al.* Prevention of injuries among male soccer players: a prospective, randomized intervention study targeting players with previous injuries or reduced function. *Am J Sports Med* 2008;36:1052–60.
- 14 Soligard T, Myklebust G, Steffen K, *et al.* Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *BMJ* 2008;337:a2469.
- 15 Hölmich P, Uhrskou P, Ulnits L, *et al.* Effectiveness of active physical training as treatment for long-standing adductor-related groin pain in athletes: randomised trial. *Lancet* 1999;353:439–43.
- 16 Machotka Z, Kumar S, Perraton LG. A systematic review of the literature on the effectiveness of exercise therapy for groin pain in athletes. *Sports Med Arthrosc Rehabil Ther Technol* 2009;1:5.
- 17 McCarthy A, Vicenzino B. Treatment of osteitis pubis via the pelvic muscles. *Man Ther* 2003;8:257–60.
- 18 Rodriguez C, Miguel A, Lima H, *et al.* Osteitis pubis syndrome in the professional soccer athlete: a case report. *J Athl Train* 2001;36:437–40.
- 19 Verrall GM, Slavotinek JP, Fon GT, *et al.* Outcome of conservative management of athletic chronic groin injury diagnosed as pubic bone stress injury. *Am J Sports Med* 2007;35:467–74.
- 20 Wollin M, Lovell G. Osteitis pubis in four young football players: a case series demonstrating successful rehabilitation. *Phys Ther Sport* 2006;7:173–4.
- 21 Aagaard P, Simonsen EB, Andersen JL, *et al.* Neural inhibition during maximal eccentric and concentric quadriceps contraction: effects of resistance training. *J Appl Physiol* 2000;89:2249–57.
- 22 Andersen LL, Andersen CH, Mortensen OS, *et al.* Muscle activation and perceived loading during rehabilitation exercises: comparison of dumbbells and elastic resistance. *Phys Ther* 2010;90:538–49.
- 23 Andersen LL, Kjaer M, Andersen CH, *et al.* Muscle activation during selected strength exercises in women with chronic neck muscle pain. *Phys Ther* 2008;88:703–11.
- 24 Andersen LL, Magnusson SP, Nielsen M, *et al.* Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Phys Ther* 2006;86:683–97.
- 25 Thorborg K, Hölmich P, Christensen R, *et al.* The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. *Br J Sports Med* 2011;45:478–91.
- 26 Hölmich P, Hölmich LR, Bjerg AM. Clinical examination of athletes with groin pain: an intraobserver and interobserver reliability study. *Br J Sports Med* 2004;38:446–51.
- 27 Jensen J, Hölmich P, Bandholm T, *et al.* Eccentric strengthening effect of hip-adductor training with elastic bands in soccer players: a randomised controlled trial. *Br J Sports Med* Published Online First: 4 July 2012. doi:10.1136/bjsports-2012-091095
- 28 Tyler TF, Nicholas SJ, Campbell RJ, *et al.* The effectiveness of a preseason exercise program to prevent adductor muscle strains in professional ice hockey players. *Am J Sports Med* 2002;30:680–3.
- 29 Sisto SA, Dyson-Hudson T. Dynamometry testing in spinal cord injury. *J Rehabil Res Dev* 2007;44:123–36.
- 30 Sundstrup E, Jakobsen MD, Andersen CH, *et al.* Muscle activation strategies during strength training with heavy loading vs. repetitions to failure. *J Strength Cond Res* 2012;26:1897–903.
- 31 Jakobsen MD, Sundstrup E, Andersen CH, *et al.* Evaluation of muscle activity during a standardized shoulder resistance training bout in novice individuals. *J Strength Cond Res* 2012;26:2515–22.
- 32 Fleck SJ, Kraemer WJ. *Designing resistance training programs*. 3rd edn. Champaign, IL: Human Kinetics, 2004.
- 33 Hermens HJ, *Commission des Communautés européennes Biomedical and Health Research Programme. SENIAM: European recommendations for surface electromyography: results of the SENIAM project*. [Pays-Bas]: Roessingh Research and Development, 1999.
- 34 Claiborne TL, Timmons MK, Pincivero DM. Test-retest reliability of cardinal plane isokinetic hip torque and EMG. *J Electromyogr Kinesiol* 2009;19:e345–352.
- 35 Aminaka N, Pietrosimone BG, Armstrong CW, *et al.* Patellofemoral pain syndrome alters neuromuscular control and kinetics during stair ambulation. *J Electromyogr Kinesiol* 2011;21:645–51.
- 36 Bolgia LA, Uhl TL. Reliability of electromyographic normalization methods for evaluating the hip musculature. *J Electromyogr Kinesiol* 2007;17:102–11.
- 37 Hodges PW, Richardson CA. Contraction of the abdominal muscles associated with movement of the lower limb. *Phys Ther* 1997;77:132–42.
- 38 Anders C, Wagner H, Puta C, *et al.* Trunk muscle activation patterns during walking at different speeds. *J Electromyogr Kinesiol* 2007;17:245–52.
- 39 Barton CJ, Coyle JA, Tinley P. The effect of heel lifts on trunk muscle activation during gait: a study of young healthy females. *J Electromyogr Kinesiol* 2009;19:598–606.
- 40 Thorborg K, Petersen J, Magnusson SP, *et al.* Clinical assessment of hip strength using a hand-held dynamometer is reliable. *Scand J Med Sci Sports* 2010;20:493–501.
- 41 Distefano LJ, Blackburn JT, Marshall SW, *et al.* Gluteal muscle activation during common therapeutic exercises. *J Orthop Sports Phys Ther* 2009;39:532–40.
- 42 Kramer JF, Vaz MD, Vandervoort AA. Reliability of isometric hip abductor torques during examiner—and belt-registered tests. *J Gerontol* 1991;46:M47–51.
- 43 Vera-Garcia FJ, Moreside JM, McGill SM. MVC techniques to normalize trunk muscle EMG in healthy women. *J Electromyogr Kinesiol* 2010;20:10–6.
- 44 Fleiss JL. *The design and analysis of clinical experiments*. New York: Wiley, 1986.
- 45 Zebis MK, Skotte J, Andersen CH, *et al.* Kettlebell swing targets semitendinosus and supine leg curl targets biceps femoris: an EMG study with rehabilitation implications. *Br J Sports Med* Published Online First: 6 July 2012. doi:10.1136/bjsports-2011-090281
- 46 Rhea MR, Alvar BA, Burkett LN, *et al.* A meta-analysis to determine the dose response for strength development. *Med Sci Sports Exerc* 2003;35:456–64.
- 47 Clark DR, Lambert MI, Hunter AM. Muscle activation in the loaded free barbell squat: a brief review. *J Strength Cond Res* 2012;26:1169–78.
- 48 Pereira GR, Leporace G, Chagas D, *et al.* Influence of hip external rotation on hip adductor and rectus femoris myoelectric activity during a dynamic parallel squat. *J Strength Cond Res* 2010;24:2749–54.
- 49 Dwyer MK, Boudreau SN, Mattacola CG, *et al.* Comparison of lower extremity kinematics and hip muscle activation during rehabilitation tasks between sexes. *J Athl Train* 2010;45:181–90.
- 50 Nicholas SJ, Tyler TF. Adductor muscle strains in sport. *Sports Med* 2002;32:339–44.
- 51 Charnock BL, Lewis CL, Garrett WE, *et al.* Adductor longus mechanics during the maximal effort soccer kick. *Sports Biomech* 2009;8:223–34.
- 52 Hysomallis C. Hip adductors' strength, flexibility, and injury risk. *J Strength Cond Res* 2009;23:1514–7.
- 53 McCully KK, Faulkner JA. Injury to skeletal muscle fibers of mice following lengthening contractions. *J Appl Physiol* 1985;59:119–26.
- 54 Armstrong RB, Ogilvie RW, Schwane JA. Eccentric exercise-induced injury to rat skeletal muscle. *J Appl Physiol* 1983;54:80–93.

- 55 Fridén J, Lieber RL. Eccentric exercise-induced injuries to contractile and cytoskeletal muscle fibre components. *Acta Physiol Scand* 2001;171:321–6.
- 56 Stauber WT, Fritz VK, Vogelbach DW, et al. Characterization of muscles injured by forced lengthening. I. Cellular infiltrates. *Med Sci Sports Exerc* 1988;20:345–53.
- 57 Tyler TF, Nicholas SJ, Campbell RJ, et al. The association of hip strength and flexibility with the incidence of adductor muscle strains in professional ice hockey players. *Am J Sports Med* 2001;29:124–8.
- 58 Petersen J, Thorborg K, Nielsen MB, et al. Preventive effect of eccentric training on acute hamstring injuries in men's soccer: a cluster-randomized controlled trial. *Am J Sports Med* 2011;39:2296–303.
- 59 Heiderscheidt BC, Hoerth DM, Chumanov ES, et al. Identifying the time of occurrence of a hamstring strain injury during treadmill running: a case study. *Clin Biomech (Bristol, Avon)* 2005;20:1072–8.
- 60 Schache AG, Wrigley TV, Baker R, et al. Biomechanical response to hamstring muscle strain injury. *Gait Posture* 2009;29:332–8.
- 61 Thorborg K, Couppé C, Petersen J, et al. Eccentric hip adduction and abduction strength in elite soccer players and matched controls: a cross-sectional study. *Br J Sports Med* 2011;45:10–3.
- 62 Thorborg K, Serner A, Petersen J, et al. Hip adduction and abduction strength profiles in elite soccer players: implications for clinical evaluation of hip adductor muscle recovery after injury. *Am J Sports Med* 2011;39:121–6.
- 63 Escamilla RF, McTaggart MSC, Fricklas EJ, et al. An electromyographic analysis of commercial and common abdominal exercises: implications for rehabilitation and training. *J Orthop Sports Phys Ther* 2006;36:45–57.
- 64 Escamilla RF, Lewis C, Bell D, et al. Core muscle activation during Swiss ball and traditional abdominal exercises. *J Orthop Sports Phys Ther* 2010;40:265–76.
- 65 Escamilla RF, Babb E, DeWitt R, et al. Electromyographic analysis of traditional and nontraditional abdominal exercises: implications for rehabilitation and training. *Phys Ther* 2006;86:656–71.
- 66 Youdas JW, Guck BR, Hebrink RC, et al. An electromyographic analysis of the Ab-Slide exercise, abdominal crunch, supine double leg thrust, and side bridge in healthy young adults: implications for rehabilitation professionals. *J Strength Cond Res* 2008;22:1939–46.
- 67 Lees A, Asai T, Andersen TB, et al. The biomechanics of kicking in soccer: a review. *J Sports Sci* 2010;28:805–17.
- 68 Cetin C, Sekir U, Yildiz Y, et al. Chronic groin pain in an amateur soccer player. *Br J Sports Med* 2004;38:223–4.
- 69 Farina D. Interpretation of the surface electromyogram in dynamic contractions. *Exerc Sport Sci Rev* 2006;34:121–7.
- 70 Farina D, Merletti R, Enoka RM. The extraction of neural strategies from the surface EMG. *J Appl Physiol* 2004;96:1486–95.
- 71 Brophy RH, Backus SI, Pansy BS, et al. Lower extremity muscle activation and alignment during the soccer instep and side-foot kicks. *J Orthop Sports Phys Ther* 2007;37:260–8.
- 72 Kaneda K, Sato D, Wakabayashi H, et al. EMG activity of hip and trunk muscles during deep-water running. *J Electromyogr Kinesiol* 2009;19:1064–70.



EMG evaluation of hip adduction exercises for soccer players: implications for exercise selection in prevention and treatment of groin injuries

Andreas Serner, Markus Due Jakobsen, Lars Louis Andersen, et al.

Br J Sports Med published online March 19, 2013

doi: 10.1136/bjsports-2012-091746

Updated information and services can be found at:
<http://bjsm.bmj.com/content/early/2013/03/18/bjsports-2012-091746.full.html>

These include:

References

This article cites 67 articles, 28 of which can be accessed free at:
<http://bjsm.bmj.com/content/early/2013/03/18/bjsports-2012-091746.full.html#ref-list-1>

P<P

Published online March 19, 2013 in advance of the print journal.

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Football \(soccer\)](#) (209 articles)

Notes

Advance online articles have been peer reviewed, accepted for publication, edited and typeset, but have not yet appeared in the paper journal. Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>